

Muiden maiden suositukset 3. annosten käyttöindikaatioista ja käytetyistä rokotteista / annoskoosta

Hanna Nohynek 6.4.2022

Terveyden ja hyvinvoinnin laitos



TECHNICAL REPORT

Interim public health considerations for the provision of additional COVID-19 vaccine doses

1 September 2021

Key messages

- Providing all eligible individuals with the recommended dose regimen should remain the current priority for COVID-19 vaccination programmes in the European Union/European Economic Area (EU/EEA).
- It is important to distinguish between 'booster' doses for people who responded adequately to primary vaccination and additional doses for those with weakened immune systems who did not respond adequately. Booster doses are given to vaccinated people (i.e. those who have completed a primary series of COVID-19 vaccination) to restore protection after it would have waned. On the other hand, additional doses as part of a primary vaccination series may be given to people with severely weakened immune systems, as they may not achieve an adequate level of protection from the standard primary vaccination.
- When assessing the need for possible booster doses of COVID-19 vaccine from the public health perspective, it is important to keep in mind the hand to be the vaccination strategy (i.e. proporting source cases of COVID 10) Vassing affectiveness against source disease should




Lokakuun alussa moni maa päättänyt aloittaa 3. annoksien annon

Ensimmäiseksi vakavasti immuunivajavaiset, sitten

- Ruotsi: hoitolaitoksessa asuvat, >80v
- Norja: kaikki 65 v täyttäneet
- Tanska:?
- UK: > 50v ja lääketieteelliset riskiryhmät









WHO SAGEn lokakuun kokouksen taustamateriaaleja viime viikolta


3.annoksetKRAR 6.4.2022

40 countries have already confirmed a COVID-19 vaccine boosters/additional doses

PLEASE NOTE: WHO DOES NOT RECOMMEND ANNUAL BOOSTER SHOTS AND HAVE CALLED FOR A TWO-MONTH VACCINE BOOSTER MORATORIUM

INDICATIVE // NON-EXHAUSTIVE

Status of COVID-19 booster / additional dose administration, # of countries



Key take-aways

- 30 countries already started administering boosters/ additional doses as of Sep. 22:
 - -2/3 of these, are HICs
 - 2 LMICs: Indonesia & Cambodia, both for HCWs only
 - UAE was the 1st
 country to start a
 booster programme in
 May 2021
- 10 HICs confirmed a booster programme but have not started it yet
- At least 11 other countries are considering a booster programme

Keille annetaan 3. koronarokoteannoksia?

Tiedot 40 eri maan osalta, tilanne 22.9.2021

- Ikäihmiset
 ikäraja vaihtelee 50+ / 60+ / 65+ / 70+ / 80+ / 86+ vuotta
- Immuunivajavaiset
- Sosiaali- ja terveydenhuollon ammattilaiset
- Sekä homologista että heterologista annostelua käytetään
- Aikaväli 2. annoksesta 3. annokseen vaihtelee riippuen kohderyhmästä

At least 30 countries are already administering booster/additional doses (1/3)

PLEASE NOTE: WHO DOES NOT RECOMMEND ANNUAL BOOSTER SHOTS AND HAVE CALLED FOR A TWO-MONTH VACCINE BOOSTER MORATORIUM

INDICATIVE // NON-EXHAUSTIVE

| | Country | Prioritized groups | | | Vx to require | Vx used | | |
|--|-----------------------|--------------------|-----------------------|------------------|-------------------------------------|--|--------|---|
| Status | | Elderly | ICP ¹ | HCW ² | booster | for booster | Start | Comment |
| Booster campaign | UAE | ✓ | / | / | Sinopharm | Pfizer/ Sinopharm | May-21 | Concerns over insufficient anti-body response related to Sinopharm. Available 6 months (Abu Dhabi) or 3 months (Dubai) after 2 nd shot |
| started | Bahrain | √ 50- | - 🗸 | / | Sinopharm | Pfizer | Jun-21 | Prioritized groups to receive booster 6 months after second shot |
| | C ∗ Turkey | √ 50+ | - | _ | Pfizer, SinoVac | Pfizer | Jun-21 | 2 nd course of mRNA vaccine available for vaccination passport |
| | | | | | | | | On Sep 10 th announced that third doses have cut hospitalization rates |
| | Dominican Republic | / | / | / | AZ, SinoVac | Pfizer/AZ | Jul-21 | Booster is voluntary. 12 weeks after 2nd shot for AZ, 4 weeks after 2nd shot for Sinovac. Pfizer not yet incl.in plan (started arriving 11 June) |
| | | ✓ 60- | - 🗸 | | Pfizer | Pfizer | Jul-21 | Gradually expanding program – since Aug 24, third shots have been available to everyone 30+. Eligible for booster 5 months after 2 nd dose |
| | Russia | / | ✓ | ✓ | Sputnik | Sputnik | Jul-21 | Russia has started booster shots for people who have been immunized more than 6 months ago |
| | Thailand | | | √ | Sinovac | AZ/Pfizer | Jul-21 | Booster shots will be administered to ~3 million people who have already received their two shots |
| | Lux- embourg | ✓ | ✓ | | Pfizer, Moderna, Janssen | Pfizer, Moderna, Janssen | Jul-21 | No plans to offer boosters for the general population; only prioritized groups |
| | Indonesia | | | ✓ | Sinovac | Moderna | Aug-21 | Currently started in Jakarta; targeted to reach 1.5 million medical workers within 2 weeks |
| | Hungary | ✓ | ✓ | | AZ, Janssen, Moderna, Pfizer, | AZ, Janssen, Moderna, Pfizer, Sinopharm, | Aug-21 | Offered to everyone 4 months after second dose; Government's official guidelines suggest a mix-and-match vaccination regimen, with final decision left to GPs |
| 1 Immunocompromised population2 Health care workforce | | | Sinopharm, Sputnik | | | No priority groups; elderly and ICP are encouraged to get booster shot | | |

At least 30 countries are already administering booster/additional doses (2/3)

PLEASE NOTE: WHO DOES NOT RECOMMEND BOOSTER SHOTS AND HAVE CALLED FOR A TWO-MONTH VACCINE BOOSTER MORATORIUM

INDICATIVE // NON-EXHAUSTIVE

| Status | Country | | Prioritized groups | | | Vx to require | Vx used | | | |
|--------------------------------|--------------|-------------|--------------------|-----|------------------|------------------|-----------------------|------------------------|--------|--|
| | | ry | Elde | rly | ICP ¹ | HCW ² | booster | for booster | Start | Comment |
| Booster campaign started | And the last | Cambodia | | | | ✓ | Sinopharm/ Sinovac | AZ/Sinovac | Aug-21 | Offered to everyone 12+ years; priorities for the boosters are medical staff, government employees and frontline workers |
| | | Lithuania | / | | ✓ | / | Not specified | Pfizer | Aug-21 | Booster given to ICP 180 days after 2 nd dose; everyone who wants a third dose would be eligible in Q4 |
| | | Serbia | / | | / | / | Not specified | Not specified | Aug-21 | Booster recommended for ICP and elderly but anyone vaccinated 6 months ago is eligible |
| | * | Chile | / | 86+ | | | Sinovac | Pfizer, AZ | Aug-21 | Administering booster shots to those already inoculated with Sinovac's COVID-19 vaccine; plan to widen to 55+ |
| | - | celand | | | | | Janssen | Pfizer, Moderna | Aug-21 | People who received Janssen vaccine will be offered one dose of Pfizer/BioNTech or Moderna vaccine after at least 8 weeks |
| | | Puerto Rico | | | ✓ | | Not specified | Not specified | Aug-21 | ICP at least 28 days after the second dose (~60k qualified residents) |
| | | Denmark | | | ✓ | | Not specified | Pfizer, Moderna | Aug-21 | Booster dose to be offered to ICP |
| | | Cuba | | | | | Soberana 2 | Soberana Plus | Aug-21 | Booster dose for Cuban two-shot Soberana 2 vaccine to increase efficiency |
| | | UK | / | 50+ | ✓ | / | Not specified | Pfizer, Moderna, AZ | Sep-21 | 50+ people, health and social care workers and younger people at risk are being offered a booster dose 6 months after final dose |
| | | USA | / | 65+ | / | ✓ | Not specified | Pfizer, Moderna | Sep-21 | Only high risk people, elderly and healthcare providers; FDA advisory group rejects COVID boosters for general population |
| | | Germany | / | | ✓ | | Not specified | Pfizer/ Moderna | Sep-21 | In addition to the elderly and ICP, Germany wants to offer the booster to people fully vaccinated with the AZ and J&J vaccines |

¹ Immunocompromised population

² Health care workforce

At least 30 countries are already administering booster/additional doses (3/3)

PLEASE NOTE: WHO DOES NOT RECOMMEND ANNUAL BOOSTER SHOTS AND HAVE CALLED FOR A TWO-MONTH VACCINE BOOSTER MORATORIUM

INDICATIVE // NON-EXHAUSTIVE

| Status | | Prioritized groups | | | Vx to require | Vx used | | |
|---------------------------|------------------------|--------------------|------------------|------------------|------------------|--------------------------|--------|--|
| | Country | Elderly | ICP ¹ | HCW ² | booster | for booster | Start | Comment |
| Booster programme started | France | √ 75+ | / | | Not specified | Pfizer/ Moderna | Sep-21 | Decision about vaccinating general population with a booster has not been made yet |
| | Malta | ✓ | / | | Not specified | TBC | Sep-21 | Only elderly in care homes; decision on whether to extend booster campaign to entire population outstanding |
| | Sweden | ✓ | / | | Not specified | Not specified | Sep-21 | Roll-out for high risk population to begin; Swedish Public Health agency expects third dose administration next year for gen population |
| | * China | √ 60+ | ✓ | √ | Sinopharm | Sinopharm, CanSinoBio | Sep-21 | Only high-risk population initiated; currently conducting trials on the need for booster shots 6 months after the beginning of the inoculation drive; also considering foreign booster |
| | * Uruguay | | | | Sinovac | Pfizer | Sep-21 | Specifically targeted at citizens who originally received Sinovac |
| | Singapore | √ 60+ | ✓ | | Not specified | Pfizer, Moderna | Sep-21 | Singapore kicked off its national Covid-19 vaccine booster programme on Sep 15 with ~3,200 seniors receiving the jabs |
| | Czech Republic | | | | Not specified | Pfizer, Moderna | Sep-21 | Option to have a booster shot will apply to everyone no earlier than eight months after final dose (four weeks for ICP) |
| | Qatar | √ 65+ | <u> </u> | / | Not specified | Pfizer, Moderna | Sep-21 | Priority is given to citizens who are highest risk of infection, despite everyone being eligible |
| | Italy mised population | √ 80+ | ✓ | \ | Not specified | Pfizer, Moderna | Sep-21 | Booster dose to be offered to prioritized groups only; at least six months after the 2 nd dose for all groups (except ICP – only 28 days) |

² Health care workforce

At least 10 other countries have confirmed to start offering booster/additional doses

INDICATIVE // NON-EXHAUSTIVE

| Status | Country | Prioritized groups | | | Vx to require | Vx used | | |
|--|-----------------------|--------------------|------------------|------------------|------------------|--------------------|----------------|--|
| | | Elderly | ICP ¹ | HCW ² | booster | for booster | Start | Comment |
| Booster programme confirmed, yet to be started | Portugal | ✓ | ✓ | | Not specified | Not specified | Sep-21 | Only offering booster shots to risk group (~100,000 people) |
| | Bermuda | √ 50+ | ✓ | ✓ | Not specified | Pfizer, Moderna | Sep-21 | Will likely follow US policy |
| | Greece | √ 60+ | ✓ | ✓ | Not specified | Pfizer, Moderna | Sep-21 | Booster dose to be offered to prioritized groups only; decision on general population outstanding |
| | Ireland | ✓ 80- | - 🗸 | ✓ | Not specified | TBC | Sep/Oct- 21 | Additionally, Boosters will be offered to residents of care facilities who are 65+; roll-out date could match flu-shot roll-out date |
| | Gibraltar | √ 50- | - | ✓ | Not specified | Pfizer | Sep/Nov- 21 | Booster vaccinations will be scheduled once the necessary supply of Pfizer BioNtech vaccine is delivered from the UK |
| | Belgium | | ✓ | | Any | Pfizer/Moderna | Sep/Nov- 21 | ICP consist of 300-400k people; eligible people to be invited automatically based on medical records |
| | Austria | √ 65- | - | ✓ | Any | Pfizer/Moderna | Oct-21 | Booster shot offered 6-9 months after first course to care home residents, 65+ yrs, high risk population and AZ/J&J recipients; 12 months for other 18+ people; ICP to start earlier with medical note |
| | Cyprus | √ 65+ | ✓ | ✓ | Not specified | Pfizer, Moderna | TBC | Booster dose to be offered to prioritized groups only |
| | Latvia | | ✓ | | Not specified | Not specified | TBC | Booster dose to be offered to ICP; decision on general population outstanding |
| | Canada sed population | | ✓ | | Not specified | Not specified | 2022 | Canada is planning for booster shots in 2022 and led negotiations with manufacturers, esp. for fast-track authorization (status from mid-April) |

¹ Immunocomprised population

Source: Government statements, press search


² Health care workforce

At least 23M booster doses have been administered globally

PLEASE NOTE: WHO DOES NOT RECOMMEND ANNUAL BOOSTER SHOTS AND HAVE CALLED FOR A TWO-MONTH VACCINE BOOSTER MORATORIUM

INDICATIVE // NON-EXHAUSTIVE

Total number of booster doses administered, #M doses



- At least 17 countries have started administering booster doses
- ~24M booster doses have been administered globally incl. ~10M in Turkey only
- Booster doses represent ~0.4% of the total number of doses administered (24M doses out of 5,950M)

The WHO stand on additional doses / booster doses

WHO's primary objective of SARS-CoV-2 vaccination is preventing hospitalizations and deaths globally

Robust evidence on waning VE against severe hospitalizations/deaths is still lacking. Current data on well maintained VE against severe disease remains encouraging while WHO notes waning VE against mild breakthrough infections

From the global perspective, increasing the first dose coverage globally will have a higher public health impact compared to increasing 3rd dose coverage in a small number of countries

Therefore, WHO proposes a **moratorium** on the use of booster doses until global vaccine coverage targets have been reached and more robust data are available on the (1) public health need for a 3rd dose and (2) booster dose performance assessments have been conducted

WHO acknowledges the need for ongoing evaluation of VE over time

WHO acknowledges that **special subpopulations** (immune compromised or older persons, for example) will require special attention

